

AMPHETAMINES		
		<i>T 1/2</i>
Biphetmaine, Obetrol	(amphetamine)	10-13HR
Strattera	(atomoxetine)	5-8 HR
Dexedrine	(dextroamphetamine)	10-13 HR
Desoxyn	(methamphetamine)	4-5 HR

ANALGESICS		
		<i>T 1/2</i>
Tramadol	(tramadol)	6-7 HR
Tramadol-ER	(tramadol-ER)	10-11 HR

ANTICOAGULANTS		
		<i>T 1/2</i>
Reo-Pro <b>**PLASMA**</b>	(abciximab)	30 MIN
Plavix	(clopidogrel)	≈6 HR
Lovenox	(enoxaparin)	≈4-7 HR
Ticlid	(ticlopidine)	13 HR
Coumadin	(warfarin)	20-60 HR

ANTIDEPRESSANTS, OTHER		
		<i>T 1/2</i>
Abilify	(aripiprazole)	75-94 HR
Wellbutrin	(bupropion)*	21 HR *12-30 HR
Buspar	(buspirone)	2-3 HR
Latuda	(lurasidone)	18-40 HR
Ludiomil	(maprotiline)	28-105 HR
Remeron	(mirtazapine)	20-40 HR
Serzone	(nefazodone)	4-8 HR
Invega	(paliperidone)	23 HR
Edronax, Vestra	(reboxitine)	
Desyrel	(trazodone)	5-9 HR
Geodon	(ziprasidone)	7 HR

ANTIPSYCHOTICS		
		<i>T 1/2</i>
Saptins	(asenapine)	24 HR
Clozaril	(clozapine)	12 HR
	<i>range</i>	4-66 HR
Inapsine	(droperidol)	2-3 HR
Haldol	(haloperidol) ORAL	14-37 HR
Lithane	(lithium)	18-36 HR
Loxitane	(loxapine)	5-19 HR
Moban	(molindone)	15 HR
Zyprexa, Zydys	(olanzapine)	30 HR
Orap	(pimozide)	111 ± 57 HR
Seroquel	(quetiapine)	~6 HR
Risperdal	(risperidone)	20 HR
Navane	(thiothixene)	34 HR

OTHER		
		<i>T 1/2</i>
Isovue 200 & 300*	(iopamidol)	≈2 HR*
≤ 1% Seizure		
*Excretion (Urine) 80% unchanged		
** Prolonged in renal failure		

MAO INHIBITORS		
		<i>T 1/2</i>
Marplan	(isocarboxazid)	11.6 HR
Nardil	(phenelzine)	11.6 HR
Deprenyl	(selegiline) ORAL	≈10 HR
Eldepryl	(selegiline) TDP	≈18.25 HR
Parnate	(tranylcypromine)	2.5 HR

PHENOTHIAZINES		
		<i>T 1/2</i>
Entrafon, Triavil, or Trilafon	(amitriptyline & perphenazine)	13-36 HR 9-12 HR
Thorazine	(chlorpromazine)	2-30 HR
	<i>biphasic</i>	
Prolixin	(fluphenazine) ORAL	≈15 HR
Compazine	(prochlorperazine)	14-22 HR
Phenergan	(promethazine)	≈15 HR
Mellaril	(thioridazine)	5-27 HR
Stelazine	(trifluorophenazine)	3-12 HR
Temaril	(trimeprazine)	~5 HR

SELECTIVE NOREPI REUPTAKE INHIBITORS (SNRIs)		
		<i>T 1/2</i>
Pristiq	(desvenlafaxine)	11 HR
Cymbalta	(duloxetine)	~12 HR
Effexor	(venlafaxine)	5-7 HR
Effexor-XR	(venlafaxine-ER)	24 HR

SELECTIVE SEROTONIN REUPTAKE INHIBITORS (SSRIs)		
		<i>T 1/2</i>
Celexa	(citalopram)	24-48 HR
Cymbalta	(duloxetine)	8-17 HR
Lexapro	(escitalopram)	27-32 HR
Prozac	(fluoxetine)	1-3 DAYS
Luvox	(fluvoxamine)	14-16 HR
	<i>elderly</i>	17-26 HR
Sarafem	(norfluoxetine)	≈9 DAY
Paxil	(paroxetine)	21 HR
Zoloft	(sertraline)	18-45 HR

TRICYCLIC ANTIDEPRESSANTS (TCAs)		
		<i>T 1/2</i>
Endep	(amitriptyline)	13-36 HR
Asendin	(amoxapine)	8 HR
	<i>metabolite</i>	30 HR
Norpramin	(desipramine)	15-24 HR
Adapin	(doxepin)	≈15 HR
	<i>metabolite: N-desmethyldoxepin</i>	≈31 HR
Janimine	(imipramine)	8-22 HR
Pamelor	(nortriptyline)	28-31 HR
Vivactil	(protriptyline)	54-92 HR
Surmontil	(trimipramine)	7-40 HR

- Most drugs cleared in FIVE (5) half-lives
- Immediate discontinuation of most antipsychotics and antidepressants is NOT recommended due to an increased risk of side effects and/or withdrawal symptoms